Finitely presented algebras defined by permutation relations of dihedral type

Ferran Cedó* Eric Jespers† Georg Klein

arXiv:1412.3707v1 [math.RA] 11 Dec 2014

Abstract

The class of finitely presented algebras over a field K with a set of generators $a_1, \ldots, a_n$ and defined by homogeneous relations of the form $a_1a_2\cdots a_n = a_{\sigma(1)}a_{\sigma(2)}\cdots a_{\sigma(n)}$, where σ runs through a subset H of the symmetric group Sym_n of degree n, is investigated. Groups H in which the cyclic group $\langle(1,2,\ldots,n)\rangle$ is a normal subgroup of index 2 are considered. Certain representations by permutations of the dihedral and semidihedral groups belong to this class of groups. A normal form for the elements of the underlying monoid $S_n(H)$ with the same presentation as the algebra is obtained. Properties of the algebra are derived, it follows that it is an automaton algebra in the sense of Ufnarovskiĭ. The universal group G_n of $S_n(H)$ is a unique product group, and it is the central localization of a cancellative subsemigroup of $S_n(H)$. This, together with previously obtained results on such semigroups and algebras, is used to show that the algebra $K[S_n(H)]$ is semiprimitive.

Keywords: semigroup ring, finitely presented, symmetric presentation, semigroup algebra, automaton algebra, regular language, primitive, Jacobson radical, semiprimitive, monoid, group
Mathematics Subject Classification: Primary: 16S15, 16S36, 20M05; Secondary: 20M25, 20M35, 16N20.

1 Introduction

We consider subgroups H of Sym_n, the symmetric group of order n, in which the cyclic subgroup $\langle(1,2,\ldots,n)\rangle$ is normal and of index 2. This class of groups includes certain representations by permutations of the dihedral and semi-dihedral groups. We study the properties of the monoid with the following presentation:

$$S_n(H) = \langle a_1, a_2, \ldots, a_n \mid a_1a_2\cdots a_n = a_{\sigma(1)}a_{\sigma(2)}\cdots a_{\sigma(n)},\ \sigma \in H\rangle.$$

In Section 2, we establish a normal form for the elements of this semigroup. The proof is an application of the diamond lemma and requires many combinatorial verifications. Using arguments by analogy, the number of actual cases to verify were kept to a minimum.

In Section 3, for a field K, we consider the algebra $K[S_n(H)]$ with the same presentation as the semigroup. From the normal form it follows that $K[S_n(H)]$ is an automaton algebra in the sense of Ufnarovskiĭ [16]. The universal group G_n of $S_n(H)$, by which we mean the group with the same

*Research partially supported by a grant of MICIIN (Spain) MTM2011-28992-C02-01.
†Research supported in part by Onderzoeksraad of Vrije Universiteit Brussel and Fonds voor Wetenschappelijk Onderzoek (Belgium).

presentation, is a unique product group. The element $z = a_1 a_2 \cdots a_n$ is central in $S_n(H)$, and $G_n = zS_n(H)\langle z\rangle^{-1}$, with $zS_n(H)$ a cancellative subsemigroup of $S_n(H)$. This is used to prove that $\mathcal{J}(K[S_n(H)]) = 0$, also using the fact that $\mathcal{J}(K[S_n(H)]) \subseteq K[zS_n(H)]$ from the reference literature. We conclude Section 3 with some results regarding prime ideals in the semigroup and in the algebra, which are related to further structural properties of the algebra.

Previous investigations of algebras of this type include the cases of H the full symmetric group [4], the alternating group of degree n [3, 5], and an abelian group [6]. In many cases the algebra turns out to be semiprimitive, with as noteworthy exception the alternating group, where the Jacobson radical being zero or not depends on the parity of n and the characteristic of the field K. The subgroups H considered in the present paper are of dihedral and quasi-dihedral type. In case the order of such groups is 2^m, there is only one other isomorphism class of groups which have the same order and which are also of nilpotency class $m-1$, the generalized quaternion group Q. In [7], we investigate $S_n(Q)$ and show that in this case more results can be obtained. Using different methods, it is shown that the semigroup $S_n(Q)$ is a two unique product semigroup if Q is given via a regular representation by permutations.

2 Monoids $S_n(H)$ defined by dihedral type relations

In this section we study $S_n(H)$ in case H contains the cyclic group $\langle(1, 2, \ldots, n)\rangle$ as a normal subgroup of index 2. Note that $n \geq 3$ and if $n = 3$, then $H = \mathrm{Sym}_3$ and $S_3(H) = S_3(\mathrm{Sym}_3)$ which is studied in [4]. Hence, throughout we assume that $n > 3$. So H is generated by $\lambda = (1, 2, \ldots, n)$ and μ, with $\mu \notin \langle\lambda\rangle$, and $\mu^2 \in \langle\lambda\rangle$. Since $\langle\lambda\rangle$ is a normal subgroup of H, we have $\mu\lambda = \lambda^k\mu$, for some integer k such that $1 \leq k < n$ and $k^2 \equiv 1 \mod n$ (so $(k, n) = 1$). For ease of notation, arithmetic will be done modulo n in the set of indices $\{1, 2, \ldots, n\}$. We have

$$\mu(i+1) = \mu\lambda(i) = \lambda^k\mu(i) = \mu(i) + k,$$

for all $i \in \{1, 2, \ldots, n\}$. So $k \neq 1$ as $\mu \notin \langle\lambda\rangle$. Thus, we obtain the following presentation

$$\begin{aligned} S_n(H) = \langle a_1, a_2, \ldots, a_n \mid\ & a_1 a_2 \cdots a_{n-1} a_n = a_i a_{i+1} a_{i+2} \cdots a_{i-2} a_{i-1} \\ & = a_i a_{i+k} a_{i+2k} \cdots a_{i-2k} a_{i-k},\ 1 \leq i \leq n\rangle. \end{aligned}$$

For simplicity, we denote $S_n(H)$ by S_n. It is clear that S_n has only trivial units. The aim of this section is to obtain a normal form for the elements of S_n. This requires tedious work. We begin by noting that

$$z = a_1 a_2 \cdots a_n$$

is a central element of S_n. Indeed, for all $1 \leq i \leq n$,

$$(a_1 a_2 \cdots a_n)a_i = a_i a_{i+1} \cdots a_{i-2} a_{i-1} a_i = a_i(a_1 a_2 \cdots a_n).$$

For every integer p, we define the following elements of length $n-2$ of S_n:

$$\begin{aligned} b_p &= a_{p+1} a_{p+2} \cdots a_{p-3} a_{p-2}, \\ c_p &= a_{p+k} a_{p+2k} \cdots a_{p-3k} a_{p-2k}. \end{aligned}$$

Lemma 2.1 *For every non-negative integer q,*

$$z b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} = z a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)}$$

and

$$zc_ic_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} = za_{i+1}b_{i+1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)},$$

for $1 \leq i \leq n$.

Proof. We shall prove this by induction on q. For $q = 0$, the statement is proved as follows.

$$\begin{aligned} zb_ia_{i-1} &= za_{i+1}a_{i+2}\cdots a_{i-2}a_{i-1} \\ &= (a_{i+k}a_{i+2k}\cdots a_{i-2k}a_{i-k}a_i)a_{i+1}a_{i+2}\cdots a_{i-2}a_{i-1} \\ &= a_{i+k}a_{i+2k}\cdots a_{i-2k}a_{i-k}z \\ &= za_{i+k}a_{i+2k}\cdots a_{i-2k}a_{i-k} \\ &= za_{i+k}c_{i+k}. \end{aligned}$$

Similarly, one obtains that $zc_ia_{i-k} = za_{i+1}b_{i+1}$. Suppose that $q \geq 1$ and the result is true for $q-1$. Since z is central, we have that

$$\begin{aligned} &zb_ib_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)} \\ &\quad = b_ib_{i-(k+1)}\cdots b_{i-(q-1)(k+1)}zb_{i-q(k+1)}a_{i-1-q(k+1)} \\ &\quad = b_ib_{i-(k+1)}\cdots b_{i-(q-1)(k+1)}za_{i+k-q(k+1)}c_{i+k-q(k+1)} \\ &\quad = zb_ib_{i-(k+1)}\cdots b_{i-(q-1)(k+1)}a_{i-1-(q-1)(k+1)}c_{i+k-q(k+1)} \\ &\quad = za_{i+k}c_{i+k}c_{i+k-(k+1)}\cdots c_{i+k-(q-1)(k+1)}c_{i+k-q(k+1)}. \end{aligned}$$

The other equality is proved similarly. ∎

Let FM_n denote the free monoid on $\{a_1, a_2, \ldots, a_n\}$ (we use the same notation for the generators of FM_n and those of S_n). Denote by $\ll$ the length-lexicographical order on FM_n generated by

$$a_1 \ll a_2 \ll \cdots \ll a_n.$$

Consider the following transformations of words in the free monoid FM_n on $\{a_1, a_2, \ldots, a_n\}$, that are dependent on the parameters i (or j), v and q, with $1 \leq i \leq n$, $v \in \mathrm{FM}_n$ and $q \geq 0$. Each transformation transforms a word into another word which is smaller in the length-lexicographical order, and this is the reason why some restrictions on i (or j) are sometimes included. The transformations are:

$$t_i(a_ia_{i+1}\cdots a_{i-2}a_{i-1}) = a_1a_2\cdots a_n, \tag{1}$$

for $i = 2, 3, \ldots, n$;

$$r_{j,m}(a_ja_1^ma_2\cdots a_n) = a_1a_2\cdots a_na_ja_1^{m-1}, \tag{2}$$

for $j = 2, 3, \ldots, n$ and $m \geq 1$;

$$h_i(a_ia_{i+k}a_{i+2k}\cdots a_{i-2k}a_{i-k}) = a_1a_2\cdots a_n; \tag{3}$$

$$\begin{aligned} &d_{i,v,q}\left(a_1\cdots a_nvb_ib_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\right) \\ &\quad = a_1\cdots a_nva_{i+k}c_{i+k}c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)}, \end{aligned} \tag{4}$$

for $n-k+1 \leq i \leq n-1$;

$$\begin{aligned} &e_{i,v,q}\left(a_1\cdots a_nvc_ic_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\right) \\ &\quad = a_1\cdots a_nva_{i+1}b_{i+1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)}, \end{aligned} \tag{5}$$

for $0 \leq i \leq n-k$;

$$\begin{aligned} & s_{i,v,q}\left(a_1\cdots a_n v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\right) \\ & \quad = a_1^2(a_2\cdots a_n)^2 v c_{i+k-(k+1)}c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)}; \end{aligned} \tag{6}$$

$$\begin{aligned} & u_{i,v,q}\left(a_1\cdots a_n v a_i c_i c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\right) \\ & \quad = a_1^2(a_2\cdots a_n)^2 v b_{i+1-(k+1)}b_{i+1-2(k+1)}\cdots b_{i+1-q(k+1)}. \end{aligned} \tag{7}$$

It follows at once from the presentation of S_n, the centrality of $z = a_1\cdots a_n$, and Lemma 2.1, that each of these transformations maps a word in FM_n to another word that represents the same element in S_n.

Let $w_1, w_2 \in \mathrm{FM}_n$. We say that w_1 covers w_2 if w_2 is obtained from w_1 by applying exactly one of the transformations $t_i, r_{j,m}, h_i, d_{i,v,q}, e_{i,v,q}, s_{i,v,q}, u_{i,v,q}$ to a subword of w_1. In this case we write $w_1 \succ w_2$. We define a partial order in FM_n by $w \geq w'$ if and only if there exist $w_0, w_1, \ldots, w_r \in \mathrm{FM}_n$, $(r \geq 0)$, such that

$$w = w_0 \succ w_1 \succ \ldots \succ w_r = w'.$$

Note that for $w, w' \in \mathrm{FM}_n$,

$$w \geq w' \Rightarrow w \gg w'.$$

Lemma 2.2 *For an integer $q \geq 0$ and any word $v \in \mathrm{FM}_n$, let*

$$\begin{cases} w_1 = a_1a_2\cdots a_n v c_i c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \in \mathrm{FM}_n \\ w_2 = a_1a_2\cdots a_n v a_{i+1}b_{i+1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \in \mathrm{FM}_n, \end{cases}$$

or

$$\begin{cases} w_1 = a_1a_2\cdots a_n v b_i b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)} \in \mathrm{FM}_n \\ w_2 = a_1a_2\cdots a_n v a_{i+k}c_{i+k}c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)} \in \mathrm{FM}_n\,. \end{cases}$$

Then there exists $w_3 \in \mathrm{FM}_n$, such that $w_1, w_2 \geq w_3$, and w_3 begins with $a_1a_2\cdots a_n$.

Proof. We will prove the first case, the second case can be proved similarly. Recall that

$$c_{i-p(k+1)} = a_{i+k-p(k+1)}a_{i+2k-p(k+1)}\cdots a_{i-3k-p(k+1)}a_{i-2k-p(k+1)}.$$

First, assume there exists an integer p such that $0 \leq p \leq q$ and $i+k-p(k+1) \equiv l \mod n$ for some $0 \leq l \leq n-k$. Let p be the smallest such number. Then

$$\begin{aligned} & e_{i+k-p(k+1),vc_i\cdots c_{i-(p-1)(k+1)},q-p}(w_1) \\ & = e_{i+k-p(k+1),vc_i\cdots c_{i-(p-1)(k+1)},q-p}(a_1\cdots a_n v c_i\cdots c_{i-(p-1)(k+1)}c_{i-p(k+1)}\cdots a_{i-k-q(k+1)}) \\ & = a_1a_2\cdots a_n v c_i\cdots c_{i-(p-1)(k+1)}a_{i+1-p(k+1)}b_{i+1-p(k+1)}\cdots b_{i+1-q(k+1)} \\ & = w_3. \end{aligned}$$

So $w_1 \geq w_3$. In case $p = 0$, clearly $w_3 = w_2$ and thus $w_2 \geq w_3$. So the result follows. In case $p \neq 0$, by the minimality condition, for all $0 \leq p' < p$ we have $i+k-p'(k+1) \equiv l \mod n$ for some $n-k+1 \leq l \leq n-1$. Since $b_ja_{j-1} = a_{j+1}b_{j+1}$ and $c_ja_{j+1-(k+1)} = a_{j+k}c_{j+k}$, we can successively apply the transformations $d_{i,v,0}$, $d_{i-(k+1),vc_i,0}$, $d_{i-2(k+1),vc_ic_{i-(k+1)},0}, \ldots, d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}$,

obtaining

$$
\begin{aligned}
&d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}d_{i,v,0}(w_2)\\
&= d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}d_{i,v,0}\\
&\quad (a_1a_2\cdots a_nva_{i+1}b_{i+1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)})\\
&= d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}d_{i,v,0}\\
&\quad (a_1a_2\cdots a_nvb_ia_{i-1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)})\\
&= d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}\\
&\quad (a_1a_2\cdots a_nva_{i+k}c_{i+k}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)})\\
&= d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}\\
&\quad (a_1a_2\cdots a_nvc_ia_{i-k}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)})\\
&= d_{i-(p-1)(k+1),vc_ic_{i-(k+1)}\cdots c_{i-(p-2)(k+1)},0}\cdots d_{i-2(k+1),vc_ic_{i-(k+1)},0}d_{i-(k+1),vc_i,0}\\
&\quad (a_1a_2\cdots a_nvc_ia_{i+1-(k+1)}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)})\\
&\vdots\\
&= a_1\cdots a_nvc_i\cdots c_{i-(p-2)(k+1)}a_{i+k-(p-1)(k+1)}c_{i+k-(p-1)(k+1)}b_{i+1-p(k+1)}\cdots b_{i+1-q(k+1)}\\
&= a_1\cdots a_nvc_i\cdots c_{i-(p-2)(k+1)}c_{i-(p-1)(k+1)}a_{i-k-(p-1)(k+1)}b_{i+1-p(k+1)}\cdots b_{i+1-q(k+1)}\\
&= a_1\cdots a_nvc_i\cdots c_{i-(p-2)(k+1)}c_{i-(p-1)(k+1)}a_{i+1-p(k+1)}b_{i+1-p(k+1)}\cdots b_{i+1-q(k+1)}\\
&= w_3.
\end{aligned}
\tag{8}
$$

So $w_2 \geq w_3$ and we have proved the statement.

Second, assume no p as above exists. One can now apply the calculation in (8) for the number $p = q + 1$ and this shows that $w_2 \geq w_3$. It it is easily seen that in this case $w_3 = w_1$ and thus the result is proved. ∎

Theorem 2.3 *Each element $a \in S_n$ can be uniquely written as a product*

$$a = a_1^i(a_2\cdots a_n)^j b, \tag{9}$$

where $b \in S_n \setminus (a_1S_n \cup a_2\cdots a_nS_n)$, i, j are non-negative integers such that if $i, j > 0$ then

$$
\begin{aligned}
b \notin \bigcup_{q\geq 0} \Big(&\bigcup_{i=n-k+1}^{n-1} S_nb_ib_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}S_n\\
&\cup \bigcup_{i=0}^{n-k} S_nc_ic_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}S_n\\
&\cup \bigcup_{i=1}^{n} \big(S_na_ib_ib_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\\
&\qquad \cup S_na_ic_ic_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}S_n\big)\Big)
\end{aligned}
$$

Proof. We will apply the diamond lemma [8, Theorem 10.4.1] to the reduction system $T = \{(w_\varphi, y_\varphi)\}$, where $w_\varphi, y_\varphi \in \mathrm{FM}_n$ are such that w_φ gets mapped to y_φ by one of the transformations (1)-(7). The partial order $\leq$ defined above is compatible with the structure of FM_n and satisfies the descending chain condition. It is clear that $y_\varphi < w_\varphi$ for every element of T, thus T is reduction-finite. To prove the result, we need to verify that all ambiguities of T are resolvable.

Let $w, w_1, w_2 \in \mathrm{FM}_n$ such that $w \succ w_1$ and $w \succ w_2$. We should show that there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2 \geq w_3$. We know that $w = a_{i_1}a_{i_2}\cdots a_{i_m}$,

$$w_1 = a_{i_1}\cdots a_{i_x}f(a_{i_{x+1}}\cdots a_{i_{x+y}})a_{i_{x+y+1}}\cdots a_{i_m}$$

and

$$w_2 = a_{i_1} \cdots a_{i_l} g(a_{i_{l+1}} \cdots a_{i_{l+p}}) a_{i_{l+p+1}} \cdots a_{i_m},$$

where $a_{i_1}, \ldots, a_{i_m} \in \{a_1, \ldots, a_n\}$ and

$$\begin{aligned} f, g \in & \{t_i \mid 2 \leq i \leq n\} \cup \{r_{j,m} \mid 2 \leq j \leq n \text{ and } m \geq 1\} \cup \{h_i \mid 1 \leq i \leq n\} \\ & \cup \{d_{i,v,q} \mid v \in \mathrm{FM}_n,\ n-k+1 \leq i \leq n-1 \text{ and } q \geq 0\} \\ & \cup \{e_{i,v,q} \mid v \in \mathrm{FM}_n,\ 0 \leq i \leq n-k \text{ and } q \geq 0\} \\ & \cup \{s_{i,v,q} \mid v \in \mathrm{FM}_n, 1 \leq i \leq n \text{ and } q \geq 0\} \\ & \cup \{u_{i,v,q} \mid v \in \mathrm{FM}_n,\ 1 \leq i \leq n \text{ and } q \geq 0\}. \end{aligned}$$

Note that if $x + y < l + 1$ or $l + p < x + 1$, then applying g to the subword $a_{i_{l+1}} \cdots a_{i_{l+p}}$ of w_1 and f to the subword $a_{i_{x+1}} \cdots a_{i_{x+y}}$ of w_2 we obtain the same word w_3 and

$$w_1 \succ w_3 \quad \text{and} \quad w_2 \succ w_3.$$

If $x + y \geq l + 1$ and $l + p \geq x + 1$, we say that the subwords $a_{i_{x+1}} \cdots a_{i_{x+y}}$ and $a_{i_{l+1}} \cdots a_{i_{l+p}}$ of w overlap. We will study all possible overlaps between subwords of the forms

(α) $a_i a_{i+1} \cdots a_{i-2} a_{i-1}$, for $2 \leq i \leq n$,

(β) $a_i a_1^m a_2 \cdots a_n$, for $2 \leq i \leq n$ and $m \geq 1$,

(δ) $a_i a_{i+k} a_{i+2k} \cdots a_{i-2k} a_{i-k}$, for $i = 1, 2, \ldots, n$,

(ζ) $a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}$,
for $v \in \mathrm{FM}_n$, $n-k+1 \leq i \leq n-1$ and $q \geq 0$,

(η) $a_1 \cdots a_n v c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)}$,
for $v \in \mathrm{FM}_n$, $0 \leq i \leq n-k$ and $q \geq 0$,

(θ) $a_1 \cdots a_n v a_i b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}$,
for $v \in \mathrm{FM}_n$, $1 \leq i \leq n$ and $q \geq 0$,

(ι) $a_1 \cdots a_n v a_i c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)}$,
for $v \in \mathrm{FM}_n$, $1 \leq i \leq n$ and $q \geq 0$.

Note that overlaps of subwords of the form (α) and (β) are dealt with in [4, Theorem 2.1]. Note also that not all kinds of subwords can overlap, and that for overlaps with the subword $a_1 \cdots a_n$ at the beginning of words of type (ζ),(η),(θ),(ι), the verification that the ambiguity is resolvable is immediate.

The remaining cases to consider are overlaps of the following form, the pairs in square brackets can be verified in a manner analogous to the pair preceding the bracket:
(α, δ), $[(\delta, \alpha)]$; (β, δ); (δ, β); (δ, δ); (ζ, α), $[(\eta, \delta)]$; (ζ, β), $[(\eta, \beta)$, (θ, β), $(\iota, \beta)]$;
(ζ, δ), $[(\eta, \alpha)]$; (θ, α), $[(\iota, \delta)]$; (θ, δ), $[(\iota, \alpha)]$;
where the notation $(\diamondsuit, \heartsuit)$ means that the ending letter(s) of a subword of the form $\diamondsuit$, overlap(s) with the beginning letter(s) of a subword of the form $\heartsuit$, as well as

(α, ζ), $[(\delta, \eta)]$; (α, η), $[(\delta, \zeta)]$; (α, θ), $[(\delta, \iota)]$; (α, ι), $[(\delta, \theta)]$;
(β, ζ); (β, η); (β, θ); (β, ι);
(ζ, ζ), $[(\eta, \eta)]$; (ζ, η), $[(\eta, \zeta)]$; (ζ, θ), $[(\eta, \iota)]$; (ζ, ι), $[(\eta, \theta)]$;
(θ, ζ), $[(\iota, \eta)]$; (θ, η), $[(\iota, \zeta)]$; (θ, θ), $[(\iota, \iota)]$; (θ, ι), $[(\iota, \theta)]$;

where the notation $(\spadesuit, \clubsuit)$ means that the ending letter(s) of a subword of the form $\spadesuit$, overlap(s) with the beginning letter(s) of a subword beginning just after the end of the subword v in a subword of the form $\clubsuit$.

Case 1: Overlaps of the form (α, δ), $[(\delta, \alpha)]$. We apply t_i to the subword of type α to get w_1 and we apply h_{i-1} to the subword of type δ to get w_2. We have

$$w = \overbrace{a_i a_{i+1} \cdots a_{i-2} \underbrace{a_{i-1}}}^{\alpha,\ t_i:\ w_1} \underbrace{a_{i-1+k} a_{i-1+2k} \cdots a_{i-1-2k} a_{i-1-k}}_{\delta,\ h_{i-1}:\ w_2}.$$

$$w_1 = a_1 a_2 \cdots a_n a_{i-1+k} a_{i-1+2k} \cdots a_{i-1-2k} a_{i-1-k}$$

and

$$w_2 = a_i a_{i+1} \cdots a_{i-2} a_1 a_2 \cdots a_n.$$

Applying a sequence of transformations r, we get

$$w_2 \ = \ a_i a_{i+1} \cdots a_{i-2} a_1 a_2 \cdots a_n \succ \ldots \succ a_1 a_2 \cdots a_n a_i a_{i+1} \cdots a_{i-2} = w_2'.$$

Then either $d_{i-1,1,0}(w_2') = w_1$ or $e_{i-1,1,0}(w_1) = w_2'$.

Case 2: Overlaps of the form (β, δ). We apply $r_{j,m}$ to the subword of type β to get w_1 and we apply h_n to the subword of type δ to get w_2. We have

$$w = \overbrace{a_j a_1^m a_2 \cdots \underbrace{a_n}}^{\beta,\ r_{j,m}:\ w_1} \underbrace{a_k a_{2k} \cdots a_{-2k} a_{-k}}_{\delta,\ h_n:\ w_2}.$$

$w_1 = a_1 \cdots a_n a_j a_1^{m-1} a_k a_{2k} \cdots a_{-2k} a_{-k}$ and $w_2 = a_j a_1^m a_2 \cdots a_{n-1} a_1 \cdots a_n$.

Applying $e_{n,a_j a_1^{m-1},0}$, we get

$$w_1 = a_1 \cdots a_n a_j a_1^{m-1} a_k a_{2k} \cdots a_{-2k} a_{-k} \succ a_1 \cdots a_n a_j a_1^{m-1} a_1 a_2 \cdots a_{-2} a_{-1}.$$

Applying a sequence of transformations r, we get

$$w_2 = a_j a_1^m a_2 \cdots a_{n-1} a_1 \cdots a_n \succ \ldots \succ a_1 \cdots a_n a_j a_1^m a_2 \cdots a_{n-1}.$$

Case 3: Overlaps of the form (δ, β). Here there are two possibilities:

(a) $a_{i-k} = a_1$. We apply h_{1+k} to the subword of type δ to get w_1 and we apply $r_{1-k,m}$ to the subword subword of type β to get w_2.

$$w = \overbrace{a_{1+k} a_{1+2k} \cdots \underbrace{a_{1-k} a_1}}^{\delta,\ h_{1+k}:\ w_1} \underbrace{a_1^{m-1} a_2 \cdots a_n}_{\beta,\ r_{1-k,m}:\ w_2}.$$

$w_1 = a_1 \cdots a_n a_1^{m-1} a_2 \cdots a_n$ and $w_2 = a_{1+k} a_{1+2k} \cdots a_{1-2k} a_1 \cdots a_n a_{1-k} a_1^{m-1}$.

If $m \neq 1$, applying a sequence of transformations r, we get

$$w_1 = a_1 \cdots a_n a_1^{m-1} a_2 \cdots a_n \succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 a_1^{m-2}.$$

Applying a sequence of transformations r, followed by h_{1+k}, followed by a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_{1+k}a_{1+2k}\cdots a_{1-2k}a_1\cdots a_n a_{1-k}a_1^{m-1} \\ &\succ \ldots \succ a_1\cdots a_n a_{1+k}a_{1+2k}\cdots a_{1-2k}a_{1-k}a_1^{m-1} \\ &\succ a_1\cdots a_n a_1\cdots a_n a_1^{m-2} \succ \ldots \succ a_1^2(a_2\cdots a_n)^2 a_1^{m-2}. \end{aligned}$$

If $m = 1$, applying a sequence of transformations r followed by $e_{1,1,0}$, we get

$$\begin{aligned} w_2 &= a_{1+k}a_{1+2k}\cdots a_{1-2k}a_1\cdots a_n a_{1-k} \succ \ldots \succ a_1\cdots a_n a_{1+k}a_{1+2k}\cdots a_{1-2k}a_{1-k} \\ &\succ a_1\cdots a_n a_2\cdots a_n = w_1. \end{aligned}$$

(b) $a_{i-k} \neq a_1$. We apply h_i to the subword of type δ to get w_1 and we apply $r_{i-k,m}$ to the subword of type β to get w_2.

$$w = \overbrace{a_i a_{i+k}\cdots a_{i-2k}a_{i-k}}^{\delta,\ h_i:\ w_1} \underbrace{a_1^m a_2\cdots a_n}_{\beta,\ r_{i-k,m}:\ w_2}.$$

$w_1 = a_1\cdots a_n a_1^m a_2\cdots a_n$ and $w_2 = a_i a_{i+k}\cdots a_{i-2k}a_1\cdots a_n a_{i-k}a_1^{m-1}$. Applying a sequence of transformations r, we get

$$w_1 = a_1\cdots a_n a_1^m a_2\cdots a_n \succ \ldots \succ a_1^2(a_2\cdots a_n)^2 a_1^{m-1}.$$

If $i \neq 1$, applying a sequence of transformations r, followed by h_i, followed by a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_i a_{i+k}\cdots a_{i-2k}a_1\cdots a_n a_{i-k}a_1^{m-1} \succ \ldots \succ a_1\cdots a_n a_i a_{i+k}\cdots a_{i-2k}a_{i-k}a_1^{m-1} \\ &\succ a_1\cdots a_n a_1\cdots a_n a_1^{m-1} \succ a_1^2(a_2\cdots a_n)^2 a_1^{m-1}. \end{aligned}$$

If $i = 1$, applying a sequence of transformations r, followed by $e_{1,1,0}$, we get

$$\begin{aligned} w_2 &= a_1 a_{1+k}\cdots a_{1-2k}a_1\cdots a_n a_{1-k}a_1^{m-1} \succ \ldots \succ a_1^2 a_2\cdots a_n a_{1+k}\cdots a_{1-2k}a_{1-k}a_1^{m-1} \\ &\succ a_1^2 a_2\cdots a_n a_2\cdots a_n a_1^{m-1}. \end{aligned}$$

Case 4: Overlaps of the form (δ, δ). We apply h_i to a subword of type δ to get w_1 and we apply $h_{i'}$ to the other subword of type δ to get w_2. We have

$$w = \overbrace{a_i a_{i+k}\cdots \underbrace{a_{i'}\cdots a_{i-k}}_{\delta,\ h_{i'}:\ w_2}\cdots a_{i'-k}}^{\delta,\ h_i:\ w_1}.$$

$w_1 = a_1\cdots a_n a_i\cdots a_{i'-k}$ and $w_2 = a_i\cdots a_{i'-k}a_1\cdots a_n$.

If $i \neq 1$, applying a sequence of transformations r, we get

$$w_2 = a_i\cdots a_{i'-k}a_1\cdots a_n \succ \ldots \succ a_1\cdots a_n a_i\cdots a_{i'-k} = w_1.$$

If $i = 1$, applying t_2, we get

$$w_1 = a_1\cdots a_n a_1 a_{1+k}\cdots a_{i'-k} \succ a_1^2 a_2\cdots a_n a_{1+k}\cdots a_{i'-k},$$

and applying a sequence of transformations r, we get

$$w_2 = a_1\cdots a_{i'-k}a_1\cdots a_n \succ \ldots \succ a_1^2 a_2\cdots a_n a_{1+k}\cdots a_{i'-k}.$$

Case 5: Overlaps of the form (ζ, α) ,$[(\eta, \delta)]$. We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $t_{i'}$ to the subword of type α to get w_2. We have

$$w = \overbrace{a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots \underbrace{a_{i'} \cdots a_{i-1-q(k+1)}}_{\alpha,\, t_{i'}:\, w_2} \cdots a_{i'-1}}^{\zeta,\, d_{i,v,q}:\, w_1}.$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_{i-q(k+1)} \cdots a_{i'-1}, \\ w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1} a_1 \cdots a_n. \end{aligned}$$

Applying $u_{i+k,v,q}$, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_{i+k-k-q(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1}. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1} a_1 \cdots a_n \\ &\succ \cdots \succ a_1^2 (a_2 \cdots a_n)^2 b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1}. \end{aligned}$$

Case 6: Overlaps of the form (ζ, β), $[(\eta, \beta)$, (θ, β), $(\iota, \beta)]$. We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $r_{j,m}$ to the subword of type β to get w_2. We have $j = i - 1 - q(k+1)$. There are two possibilities.

(a) $j = i - 1 - q(k+1) \neq 1$.

$$w = \overbrace{a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} \underbrace{a_{i-1-q(k+1)}}_{\beta,\, r_{j,m}:\, w_2}}^{\zeta,\, d_{i,v,q}:\, w_1} a_1^m a_2 \cdots a_n.$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_1^m a_2 \cdots a_n, \\ w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_1 a_2 \cdots a_n a_{i-1-q(k+1)} a_1^{m-1}. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_1^m a_2 \cdots a_n \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_1^{m-1}. \end{aligned}$$

Applying a sequence of transformations r and $d_{i,a_2 \cdots a_n v,q}$, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_1 a_2 \cdots a_n a_{i-1-q(k+1)} a_1^{m-1} \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} a_1^{m-1} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_1^{m-1}. \end{aligned}$$

(b) $j = i - 1 - q(k+1) = 1$.

$$w = \overbrace{a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} \underbrace{a_1}_{\beta,\, r_{j,m}:\, w_2}}^{\zeta,\, d_{i,v,q}:\, w_1} a_1^{m-1} a_2 \cdots a_n.$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_1^{m-1} a_2 \cdots a_n, \\ w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i-3-q(k+1)} \cdot \\ &\qquad \cdot a_1 a_2 \cdots a_n a_{i-2-q(k+1)} a_1^{m-1}. \end{aligned}$$

If $m \neq 1$, this is solved in the same way as (a). If $m = 1$, we use the facts that $a_{i+k-k-q(k+1)} = a_2$ and $a_3 \cdots a_n = a_{i+1-q(k+1)} \cdots a_{i-2-q(k+1)} = b_{i-q(k+1)}$ and applying $u_{i+k,v,q}$, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_2 \cdots a_n \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v b_{i+k+1-(k+1)} b_{i+k+1-2(k+1)} \cdots b_{i+k+1-q(k+1)} a_3 \cdots a_n \\ &= a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} b_{i-q(k+1)}. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i-3-q(k+1)} \cdot \\ &\qquad \cdot a_1 a_2 \cdots a_n a_{i-2-q(k+1)} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i-3-q(k+1)} a_{i-2-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} b_{i-q(k+1)}. \end{aligned}$$

Case 7: Overlaps of the form (ζ, δ), $[(\eta, \alpha)]$. We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $h_{i-1-q(k+1)}$ to the subword of type δ to get w_2. We have

$$w = \overbrace{a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} \underbrace{a_{i-1-q(k+1)}}_{\delta,\ h_{i-1-q(k+1)}:\ w_2}}^{\zeta,\ d_{i,v,q}:\ w_1} a_{i-1-q(k+1)+k} \cdots a_{i-1-q(k+1)-k}.$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_{i-1-q(k+1)+k} \cdots a_{i-1-q(k+1)-k}, \\ w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_1 \cdots a_n. \end{aligned}$$

Applying $u_{i+k,v,q+1}$, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} a_{i+k+k-q(k+1)-k-1} \cdots \\ &\qquad \cdots a_{i+k-k-q(k+1)-k-1} \\ &= a_1 \cdots a_n v a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} c_{i+k-(q+1)(k+1)} a_{i+k-k-(q+1)(k+1)} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)}. \end{aligned}$$

Applying a sequence of transformations r we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_1 \cdots a_n \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)}. \end{aligned}$$

Case 8: Overlaps of the form (θ, α), $[(\iota, \delta)]$. We apply $s_{i,v,q}$ to the subword of type θ to get w_1, and we apply $t_{i'}$ to the subword of type α to get w_2. We have

$$w = \overbrace{a_1 \cdots a_n v a_i b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots \underbrace{a_{i'} \cdots a_{i-1-q(k+1)}}_{\alpha,\ t_{i'}:\ w_2}}^{\theta,\ s_{i,v,q}:\ w_1} \cdots \underbrace{a_{i'-1}}_{\alpha,\ t_{i'}:\ w_2}.$$

$$\begin{aligned} w_1 &= a_1^2 (a_2 \cdots a_n)^2 v c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} a_{i-q(k+1)} \cdots a_{i'-1}, \\ w_2 &= a_1 \cdots a_n v a_i b_i b_{i-(k+1)} \cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)} \cdots a_{i'-1} a_1 \cdots a_n. \end{aligned}$$

We have

$$\begin{aligned} w_1 &= a_1^2 (a_2 \cdots a_n)^2 v c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} a_{i-q(k+1)} \cdots a_{i'-1} \\ &= a_1^2 (a_2 \cdots a_n)^2 v c_{i-1} c_{i-1-(k+1)} \cdots c_{i-1-(q-1)(k+1)} a_{i-1-k-(q-1)(k+1)} \cdots a_{i'-1}. \end{aligned}$$

Applying a sequence of transformations r, we get

$$w_2 \succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v a_i b_i b_{i-(k+1)}\cdots b_{i-(q-1)(k+1)} a_{i+1-q(k+1)}\cdots a_{i'-1} = w_2'.$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 9: Overlaps of the form (θ, δ), $[(\iota, \alpha)]$. We apply $s_{i,v,q}$ to the subword of type θ to get w_1, and we apply $h_{i-1-q(k+1)}$ to the subword of type δ to get w_2. We have

$$w = \overbrace{a_1\cdots a_n v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} \underbrace{a_{i-1-q(k+1)}}_{\delta,\, h_{i-1-q(k+1)}:\, w_2}}^{\theta,\, s_{i,v,q}:\, w_1} a_{i-1-q(k+1)+k}\cdots a_{i-1-q(k+1)-k}.$$

$$w_1 = a_1^2(a_2\cdots a_n)^2 v c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} a_{i-1-q(k+1)+k}\cdots a_{i-1-q(k+1)-k},$$
$$w_2 = a_1\cdots a_n v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} a_1\cdots a_n.$$

We have

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} a_{i-1-q(k+1)+k}\cdots a_{i-1-q(k+1)-k} \\ &= a_1^2(a_2\cdots a_n)^2 v c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-(q+1)(k+1)} a_{i+k-k-(q+1)(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v c_{i-1} c_{i-1-(k+1)}\cdots c_{i-1-q(k+1)} a_{i-1-k-q(k+1)}. \end{aligned}$$

Applying a sequence of transformations r, we get

$$w_2 \succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} = w_2'.$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 10: Overlaps of the form (α, ζ), $[(\delta, \eta)]$. We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $t_{i'}$ to the subword of type α to get w_2. There are two kinds of overlap, the first kind is as follows:

$$\begin{aligned} w &= a_1\cdots a_n v b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= \overbrace{a_1\cdots a_n v' \underbrace{a_{i'}\cdots a_{i+1}}_{\alpha,\, t_{i'}:\, w_2}\cdots a_{i'-1}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\zeta,\, d_{i,v,q}:\, w_1}. \end{aligned}$$

$$w_1 = a_1\cdots a_n v' a_{i'}\cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)},$$
$$w_2 = a_1\cdots a_n v' a_1\cdots a_n a_{i'}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}.$$

Applying h_i followed by a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1\cdots a_n v' a_{i'}\cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)} \\ &\succ a_1\cdots a_n v' a_{i'}\cdots a_{i-1} a_1\cdots a_n c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} \\ &\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-1} c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-1} c_{i-1} c_{i-1-(k+1)}\cdots c_{i-1-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_{i'}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &\succ a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-2} b_{i-k-1} b_{i-k-1-(k+1)}\cdots \\ &\qquad\qquad \cdots b_{i-k-1-(q-1)(k+1)} a_{i-k-1-1-(q-1)(k+1)} \\ &= w_2'. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

The second kind of overlap is as follows:

$$\begin{aligned} w &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= \overbrace{a_1 \cdots \underbrace{a_{i'} \cdots a_n \cdots a_{i+1} \cdots a_{i'-1}}_{\alpha,\, t_{i'}:\, w_2} \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\zeta,\, d_{i,v,q}:\, w_1}. \end{aligned}$$

Here $v = a_{n+1} \cdots a_i$. Thus

$$\begin{aligned} w_1 &= a_1 \cdots a_{i'} \cdots a_n \cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)}, \\ w_2 &= a_1 \cdots a_{i'-1} a_1 a_2 \cdots a_n a_{i'} \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}. \end{aligned}$$

Applying h_i followed by a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_{i'} \cdots a_n \cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} \\ &\succ a_1 \cdots a_{i'} \cdots a_n \cdots a_{i-1} a_1 \cdots a_n c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 a_1 \cdots a_{i-1} c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 a_1 \cdots a_{i-1} c_{i-1} c_{i-1-(k+1)} \cdots c_{i-1-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_{i'-1} a_1 a_2 \cdots a_n a_{i'} \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 a_1 \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 a_1 \cdots a_{i-2} b_{i-k-1} b_{i-k-1-(k+1)} \cdots \\ &\qquad \cdots b_{i-k-1-(q-1)(k+1)} a_{i-k-1-1-(q-1)(k+1)} \\ &= w_2'. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

Case 11: Overlaps of the form (α, η), $[(\delta, \zeta)]$. We apply $e_{i,v,q}$ to the subword of type η to get w_1 and we apply t_{i+k+1} to the subword of type α to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1 \cdots a_n v c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \\ &= \overbrace{a_1 \cdots a_n v' \underbrace{a_{i+k+1} a_{i+k+2} \cdots a_{i+k-1} a_{i+k}}_{\alpha,\, t_{i+k+1}:\, w_2} a_{i+2k} \cdots a_{i-3k} a_{i-2k} c_{i-(k+1)} c_{i-2(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)}}^{\eta,\, e_{i,v,q}:\, w_1}. \end{aligned}$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_{i+k+1} a_{i+k+2} \cdots a_{i+k-1} a_{i+1} b_{i+1} b_{i+1-(k+1)} \cdots b_{i+1-q(k+1)}, \\ w_2 &= a_1 \cdots a_n v' a_1 \cdots a_n a_{i+2k} \cdots a_{i-3k} a_{i-2k} c_{i-(k+1)} c_{i-2(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)}. \end{aligned}$$

Applying $s_{i+k+1,v',1}$, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_{i+k+1} a_{i+k+2} \cdots a_{i+k-1} a_{i+1} b_{i+1} b_{i+1-(k+1)} \cdots b_{i+1-q(k+1)} \\ &= a_1 \cdots a_n v' a_{i+k+1} b_{i+k+1} a_{i+1} b_{i+1} b_{i+1-(k+1)} b_{i+1-2(k+1)} \cdots b_{i+1-q(k+1)} \\ &= a_1 \cdots a_n v' a_{i+k+1} b_{i+k+1} b_{i+k+1-(k+1)} a_{i-1} b_{i+1-(k+1)} b_{i+1-2(k+1)} \cdots b_{i+1-q(k+1)} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v' c_{i+k} b_{i+1-(k+1)} b_{i+1-2(k+1)} \cdots b_{i+1-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 v' a_{i+2k} \cdots a_{i-2k} a_{i-k} b_{i+1-(k+1)} b_{i+1-2(k+1)} \cdots b_{i+1-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 v' a_{i+2k} \cdots a_{i-2k} a_{i-k} b_{i-k} b_{i-k-(k+1)} \cdots b_{i-k-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned}
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_{i+2k}\cdots a_{i-3k}a_{i-2k}c_{i-(k+1)}c_{i-2(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\\
&\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' a_{i+2k}\cdots a_{i-3k}a_{i-2k}c_{i-(k+1)}c_{i-2(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_{i+2k}\cdots a_{i-3k}a_{i-2k}c_{i-k-1}c_{i-k-1-(k+1)}\cdots\\
&\qquad\qquad \cdots c_{i-k-1-(q-1)(k+1)}a_{i-k-1-k-(q-1)(k+1)}\\
&= w_2'.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

Case 12: Overlaps of the form (α, θ), $[(\delta, \iota)]$. We apply $s_{i,v,q}$ to the subword of type θ to get w_1 and we apply $t_{i'}$ to the subword of type α to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}
w &= a_1\cdots a_n v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\\
&= \overbrace{a_1\cdots a_n v' \underbrace{a_{i'}\cdots a_i\cdots a_{i'-1}}_{\alpha,\, t_{i'}:\, w_2}\cdots a_{i-2}b_{i-(k+1)}b_{i-2(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}}^{\theta,\, s_{i,v,q}:\, w_1}.
\end{aligned}$$

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-1}c_{i+k-(k+1)}c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)},\\
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_{i'}\cdots a_{i-2}b_{i-(k+1)}b_{i-2(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}.
\end{aligned}$$

We have

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-1}c_{i+k-(k+1)}c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-1}c_{i-1}c_{i-1-(k+1)}\cdots c_{i-1-(q-1)(k+1)}.
\end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned}
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_{i'}\cdots a_{i-2}b_{i-(k+1)}b_{i-2(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\\
&\succ a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-2}b_{i-(k+1)}b_{i-2(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_{i'}\cdots a_{i-2}b_{i-k-1}b_{i-k-1-(k+1)}\cdots\\
&\qquad\qquad \cdots b_{i-k-1-(q-1)(k+1)}a_{i-k-1-1-(q-1)(k+1)}\\
&= w_2'.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 13: Overlaps of the form (α, ι), $[(\delta, \theta)]$. We apply $u_{i,v,q}$ to the subword of type ι to get w_1 and we apply t_{i+1} to the subword of type α to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}
w &= a_1\cdots a_n v a_i c_i c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\\
&= \overbrace{a_1\cdots a_n v' \underbrace{a_{i+1}a_{i+2}\cdots a_{i-1}a_i}_{\alpha,\, t_{i+1}:\, w_2} c_i c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}}^{\iota,\, u_{i,v,q}:\, w_1}.
\end{aligned}$$

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i+1}a_{i+2}\cdots a_{i-1}b_{i+1-(k+1)}b_{i+1-2(k+1)}\cdots b_{i+1-q(k+1)},\\
w_2 &= a_1\cdots a_n v' a_1\cdots a_n c_i c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}.
\end{aligned}$$

We have

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i+1}a_{i+2}\cdots a_{i-1}b_{i+1-(k+1)}b_{i+1-2(k+1)}\cdots b_{i+1-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_{i+1}b_{i+1}b_{i+1-(k+1)}b_{i+1-2(k+1)}\cdots b_{i+1-q(k+1)}.
\end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v' a_1 \cdots a_n c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v' c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \\ &= w_2'. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 14: Overlaps of the form (β, ζ). We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $r_{j,m}$ to the subword of type β to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1 \cdots a_n v b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= \overbrace{a_1 \cdots a_n v' \underbrace{a_j a_1^{m-1} a_1 \cdots a_{i+1} \cdots a_n}_{\beta,\, r_{j,m}:\, w_2} \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\zeta,\, d_{i,v,q}:\, w_1}. \end{aligned}$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_j a_1^{m-1} a_1 \cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)}, \\ w_2 &= a_1 \cdots a_n v' a_1 \cdots a_n a_j a_1^{m-1} a_1 \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}. \end{aligned}$$

Applying h_i and a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_j a_1^{m-1} a_1 \cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} \\ &\succ a_1 \cdots a_n v' a_j a_1^{m-1} a_1 \cdots a_{i-1} a_1 a_2 \cdots a_n c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} \\ &\succ \ldots \succ a_1^2 (a_2 \cdots a_n)^2 v' a_j a_1^{m-1} a_1 \cdots a_{i-1} c_{i+k-(k+1)} c_{i+k-2(k+1)} \cdots c_{i+k-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 v' a_j a_1^{m-1} a_1 \cdots a_{i-1} c_{i-1} c_{i-1-(k+1)} \cdots c_{i-1-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v' a_1 \cdots a_n a_j a_1^{m-1} a_1 \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &\succ a_1^2 (a_2 \cdots a_n)^2 v' a_j a_1^{m-1} a_1 \cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= a_1^2 (a_2 \cdots a_n)^2 v' a_j a_1^{m-1} a_1 \cdots a_{i-2} b_{i-k-1} b_{i-k-1-(k+1)} \cdots \\ &\qquad \cdots b_{i-k-1-(q-1)(k+1)} a_{i-k-1-1-(q-1)(k+1)} \\ &= w_2'. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

Case 15: Overlaps of the form (β, η). We apply $e_{n-k,v,q}$ to the subword of type η to get w_1 and we apply $r_{j,m}$ to the subword of type β to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1 \cdots a_n v c_{n-k} c_{n-k-(k+1)} \cdots c_{n-k-q(k+1)} a_{n-k-k-q(k+1)} \\ &= \overbrace{a_1 \cdots a_n v' \underbrace{a_j a_1^{m-1} a_1 \cdots a_{n-1} a_n}_{\beta,\, r_{j,m}:\, w_2} a_{n+k} \cdots a_{n-3k} c_{n-k-(k+1)} c_{n-k-2(k+1)} \cdots c_{n-k-q(k+1)} a_{n-k-k-q(k+1)}}^{\eta,\, e_{n-k,v,q}:\, w_1}. \end{aligned}$$

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_j a_1^{m-1} a_1 \cdots a_{n-1} a_{n-k+1} b_{n-k+1} b_{n-k+1-(k+1)} \cdots b_{n-k+1-q(k+1)}, \\ w_2 &= a_1 \cdots a_n v' a_1 \cdots a_n a_j a_1^{m-1} a_{n+k} \cdots a_{n-3k} c_{n-k-(k+1)} c_{n-k-2(k+1)} \cdots \\ &\qquad \cdots c_{n-k-q(k+1)} a_{n-k-k-q(k+1)}. \end{aligned}$$

Applying $s_{1,v'a_ja_1^{m-1},1}$, we get

$$\begin{aligned}
w_1 &= a_1\cdots a_n v' a_j a_1^{m-1} a_1\cdots a_{n-1} a_{n-k+1} b_{n-k+1} b_{n-k+1-(k+1)}\cdots b_{n-k+1-q(k+1)}\\
&= a_1\cdots a_n v' a_j a_1^{m-1} a_1 b_1 b_{1-(k+1)} a_{n-k-1} b_{n-k+1-(k+1)} b_{n-k+1-2(k+1)}\cdots b_{n-k+1-q(k+1)}\\
&\succ a_1^2(a_1\cdots a_n)^2 v' a_j a_1^{m-1} c_{1+k-(k+1)} b_{n-k+1-(k+1)} b_{n-k+1-2(k+1)}\cdots b_{n-k+1-q(k+1)}\\
&= a_1^2(a_1\cdots a_n)^2 v' a_j a_1^{m-1} a_k\cdots a_{-2k} b_{n-k+1-(k+1)} b_{n-k+1-2(k+1)}\cdots b_{n-k+1-q(k+1)}\\
&= a_1^2(a_1\cdots a_n)^2 v' a_j a_1^{m-1} a_k\cdots a_{-2k} b_{-2k} b_{-2k-(k+1)}\cdots b_{-2k-(q-1)(k+1)}\\
&= w_1'.
\end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned}
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_j a_1^{m-1} a_{n+k}\cdots a_{n-3k} c_{n-k-(k+1)} c_{n-k-2(k+1)}\cdots\\
&\qquad\qquad \cdots c_{n-k-q(k+1)} a_{n-k-k-q(k+1)}\\
&\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_{n+k}\cdots a_{n-3k} c_{n-k-(k+1)} c_{n-k-2(k+1)}\cdots\\
&\qquad\qquad \cdots c_{n-k-q(k+1)} a_{n-k-k-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_{n+k}\cdots a_{n-3k} c_{n-2k-1} c_{n-2k-1-(k+1)}\cdots\\
&\qquad\qquad \cdots c_{n-2k-1-(q-1)(k+1)} a_{n-2k-1-k-(q-1)(k+1)}\\
&= w_2'.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

Case 16: Overlaps of the form (β, θ). We apply $s_{i,v,q}$ to the subword of type θ to get w_1 and we apply $r_{j,m}$ to the subword of type β to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}
w &= a_1\cdots a_n v a_i b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\
&= \overbrace{a_1\cdots a_n v' \underbrace{a_j a_1^{m-1} a_1\cdots a_i\cdots a_n}_{\beta,\ r_{j,m}:\ w_2}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\theta,\ s_{i,v,q}:\ w_1}.
\end{aligned}$$

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1\cdots a_{i-1} c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)},\\
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_j a_1^{m-1} a_{n+1}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}.
\end{aligned}$$

We have

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1\cdots a_{i-1} c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1\cdots a_{i-1} c_{i-1} c_{i-1-(k+1)}\cdots c_{i-1-(q-1)(k+1)}.
\end{aligned}$$

Applying a sequence of transformations r, we get

$$\begin{aligned}
w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_j a_1^{m-1} a_{n+1}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}.\\
&\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_{n+1}\cdots a_{i-2} b_{i-(k+1)} b_{i-2(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\
&= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_{n+1}\cdots a_{i-2} b_{i-k-1} b_{i-k-1-(k+1)}\cdots\\
&\qquad\qquad \cdots b_{i-k-1-(q-1)(k+1)} a_{i-k-1-1-(q-1)(k+1)}\\
&= w_2'.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 17: Overlaps of the form (β, ι). We apply $u_{n,v,q}$ to the subword of type ι to get w_1 and we

apply $r_{j,m}$ to the subword of type β to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1\cdots a_n v a_n c_n c_{n-(k+1)}\cdots c_{n-q(k+1)}a_{n-k-q(k+1)} \\ &= \overbrace{a_1\cdots a_n v' \underbrace{a_j a_1^{m-1} a_1\cdots a_n}_{\beta,\, r_{j,m}:\, w_2} c_n c_{n-(k+1)}\cdots c_{n-q(k+1)}a_{n-k-q(k+1)}}^{\iota,\, u_{n,v,q}:\, w_1}. \end{aligned}$$

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1\cdots a_{n-1} b_{n+1-(k+1)} b_{n+1-2(k+1)}\cdots b_{n+1-q(k+1)} \\ w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_j a_1^{m-1} c_n c_{n-(k+1)}\cdots c_{n-q(k+1)} a_{n-k-q(k+1)}. \end{aligned}$$

We have

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1\cdots a_{n-1} b_{n+1-(k+1)} b_{n+1-2(k+1)}\cdots b_{n+1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1 b_{n+1} b_{n+1-(k+1)} b_{n+1-2(k+1)}\cdots b_{n+1-q(k+1)}. \end{aligned}$$

Applying a sequence of transformations r followed by $e_{n,a_2\cdots a_n v' a_j a_1^{m-1},q}$, we get

$$\begin{aligned} w_2 &= a_1\cdots a_n v' a_1\cdots a_n a_j a_1^{m-1} c_n c_{n-(k+1)}\cdots c_{n-q(k+1)} a_{n-k-q(k+1)} \\ &\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} c_n c_{n-(k+1)}\cdots c_{n-q(k+1)} a_{n-k-q(k+1)} \\ &\succ a_1^2(a_2\cdots a_n)^2 v' a_j a_1^{m-1} a_1 b_{n+1} b_{n+1-(k+1)} b_{n+1-2(k+1)}\cdots b_{n+1-q(k+1)} \\ &= w_1. \end{aligned}$$

Case 18: Overlaps of the form (ζ,ζ), $[(\eta,\eta)]$. We apply $d_{i,v,q}$ to a subword of type ζ to get w_1, and we apply $d_{i',v',q'}$ to the other subword of type ζ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1\cdots a_n v b_i b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= \overbrace{\underbrace{a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i+1}\cdots a_{i'-1-q'(k+1)}}_{\zeta,\, d_{i',v',q'}:\, w_2}\cdots a_{i-2} b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\zeta,\, d_{i,v,q}:\, w_1}. \end{aligned}$$

Without loss of generality, it is assumed that the subwords $a_1\cdots a_n$ at the beginning of the two words of type ζ coincide. We have

$$\begin{aligned} w_1 &= a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)}, \\ w_2 &= a_1\cdots a_n v' a_{i'+k} c_{i'+k} c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)}\cdot \\ &\qquad \cdot a_{i'-q'(k+1)}\cdots a_{i-2} b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}. \end{aligned}$$

Applying h_i and a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_i a_{i+k} c_{i+k} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)} \\ &= a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_i a_{i+k}\cdots a_{i-k}\cdot \\ &\qquad \cdot c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} \\ &\succ a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1} a_1\cdots a_n\cdot \\ &\qquad \cdot c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} \\ &\succ \ldots \succ a_1^2(a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1}\cdot \\ &\qquad \cdot c_{i+k-(k+1)} c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1}\cdot \\ &\qquad \cdot c_{i-1} c_{i-1-(k+1)}\cdots c_{i-1-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

Applying $u_{i'+k,v',q'}$, we get

$$\begin{aligned}
w_2 &= a_1\cdots a_n v' a_{i'+k} c_{i'+k} c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)} a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\
&\qquad\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\
&= a_1\cdots a_n v' a_{i'+k} c_{i'+k} c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)} a_{i'+k-k-q'(k+1)}\cdots a_{i-2}\cdot \\
&\qquad\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\
&\succ a_1^2 (a_2\cdots a_n)^2 v' b_{i'+k+1-(k+1)}\cdots b_{i'+k+1-q'(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-2}\cdot \\
&\qquad\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\
&= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-2}\cdot \\
&\qquad\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\
&= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-2}\cdot \\
&\qquad\qquad \cdot b_{i-k-1}\cdots b_{i-k-1-(q-1)(k+1)} a_{i-k-1-1-(q-1)(k+1)} \\
&= w_2'.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2' \geq w_3$.

Case 19: Overlaps of the form (ζ, η), $[(\eta, \zeta)]$. We apply $e_{i,v,q}$ to the subword of type η to get w_1 and we apply $d_{i+(q'+1)(k+1),v',q'}$ to the subword of type ζ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}
w &= a_1\cdots a_n v c_i c_{i-(k+1)}\cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \\
&= \overbrace{\underbrace{a_1\cdots a_n v' b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)} a_{i+k}}_{\zeta,\ d_{i+(q'+1)(k+1),v',q'}:\ w_2}\cdots a_{i-2k} c_{i-(k+1)}\cdots c_{i-q(k+1)} a_{i-k-q(k+1)}}^{\eta,\ e_{i,v,q}:\ w_1}.
\end{aligned}$$

Without loss of generality, it is assumed that the subwords $a_1\cdots a_n$ at the beginning of the word of type η and at the beginning of the word of type ζ coincide. We have

$$\begin{aligned}
w_1 &= a_1\cdots a_n v' b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)} a_{i+1} b_{i+1}\cdots b_{i+1-q(k+1)}, \\
w_2 &= a_1\cdots a_n v' a_{i+(q'+1)(k+1)+k} c_{i+(q'+1)(k+1)+k}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}\cdot \\
&\qquad\qquad \cdot a_{i+2k}\cdots a_{i-2k} c_{i-(k+1)}\cdots c_{i-q(k+1)} a_{i-k-q(k+1)}.
\end{aligned}$$

Applying $d_{i+(q'+1)(k+1),v',q'+1}$, we get

$$\begin{aligned}
w_1 &= a_1\cdots a_n v' b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)} a_{i+1} b_{i+1}\cdots b_{i+1-q(k+1)} \\
&= a_1\cdots a_n v' b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)} a_{i+1} a_{i+2}\cdots a_{i-1}\cdot \\
&\qquad\qquad \cdot b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\
&= a_1\cdots a_n v' b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)} b_{i+(q'+1)(k+1)-(q'+1)(k+1)} a_{i-1}\cdot \\
&\qquad\qquad \cdot b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\
&\succ a_1\cdots a_n v' a_{i+(q'+1)(k+1)+k} c_{i+(q'+1)(k+1)+k}\cdots c_{i+(q'+1)(k+1)+k-(q'+1)(k+1)}\cdot \\
&\qquad\qquad \cdot b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\
&= a_1\cdots a_n v' a_{i+(q'+1)(k+1)+k} c_{i+(q'+1)(k+1)+k}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}\cdot \\
&\qquad\qquad \cdot a_{i+2k}\cdots a_{i-2k} a_{i-k} b_{i-k}\cdots b_{i-k-(q-1)(k+1)} \\
&= w_1'.
\end{aligned}$$

We have

$$\begin{aligned}w_2 &= a_1\cdots a_n v' a_{i+(q'+1)(k+1)+k} c_{i+(q'+1)(k+1)+k}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}\cdot\\ &\qquad\cdot a_{i+2k}\cdots a_{i-2k} c_{i-(k+1)}\cdots c_{i-q(k+1)} a_{i-k-q(k+1)}\\ &= a_1\cdots a_n v' a_{i+(q'+1)(k+1)+k} c_{i+(q'+1)(k+1)+k}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}\cdot\\ &\qquad\cdot a_{i+2k}\cdots a_{i-2k} c_{i-k-1}\cdots c_{i-k-1-(q-1)(k+1)} a_{i-k-q(k+1)}.\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2 \geq w_3$.

Case 20: Overlaps of the form (ζ, θ), $[(\eta, \iota)]$. We apply $s_{i,v,q}$ to the subword of type θ to get w_1 and we apply $d_{i',v',q'}$ to the subword of type ζ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}w &= a_1\cdots a_n v a_i b_i\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\ &= \overbrace{\underbrace{a_1\cdots a_n v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_i\cdots a_{i'-1-q'(k+1)}}_{\zeta,\, d_{i',v',q'}:\, w_2}\cdots a_{i-2} b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\theta,\, s_{i,v,q}:\, w_1}.\end{aligned}$$

Without loss of generality, it is assumed that the subwords $a_1\cdots a_n$ at the beginning of the word of type θ and at the beginning of the word of type ζ coincide. We have

$$\begin{aligned}w_1 &= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)},\\ w_2 &= a_1\cdots a_n v' a_{i'+k} c_{i'+k}\cdots c_{i'+k-q'(k+1)} a_{i'-q'(k+1)}\cdots a_{i-2}\cdot\\ &\qquad\cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}.\end{aligned}$$

We have

$$\begin{aligned}w_1 &= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1} c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)}\\ &= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-1} c_{i-1}\cdots c_{i-1-(q-1)(k+1)}.\end{aligned}$$

Applying $u_{i'+k,v',q'}$, we get

$$\begin{aligned}w_2 &= a_1\cdots a_n v' a_{i'+k} c_{i'+k}\cdots c_{i'+k-q'(k+1)} a_{i'-q'(k+1)}\cdots a_{i-2}\cdot\\ &\qquad\cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\ &= a_1\cdots a_n v' a_{i'+k} c_{i'+k}\cdots c_{i'+k-q'(k+1)} a_{i'+k-k-q'(k+1)}\cdots a_{i-2}\cdot\\ &\qquad\cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\ &\succ a_1^2 (a_2\cdots a_n)^2 v' b_{i'+k+1-(k+1)}\cdots b_{i'+k+1-q'(k+1)} a_{i'+1-q'(k+1)}\cdots\cdot a_{i-2}\cdot\\ &\qquad\cdot b_{i-(k+1)}\cdots b_{i-q(k+1)} a_{i-1-q(k+1)}\\ &= a_1^2 (a_2\cdots a_n)^2 v' b_{i'}\cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)}\cdots a_{i-2}\cdot\\ &\qquad\cdot b_{i-k-1}\cdots b_{i-k-1-(q-1)(k+1)} a_{i-1-q(k+1)}\\ &= w_2'.\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2' \geq w_3$.

Case 21: Overlaps of the form (ζ, ι), $[(\eta, \theta)]$. We apply $u_{i,v,q}$ to the subword of type ι to get w_1 and we apply $d_{i+q'(k+1)+1,v',q'}$ to the subword of type ζ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned}w &= a_1\cdots a_n v a_i c_i\cdots c_{i-q(k+1)} a_{i-k-q(k+1)}\\ &= \overbrace{\underbrace{a_1\cdots a_n v' b_{i+q'(k+1)+1}\cdots b_{i+q'(k+1)+1-q'(k+1)} a_i}_{\zeta,\, d_{i+q'(k+1)+1,v',q'}:\, w_2} c_i c_{i-(k+1)}\cdots c_{i-q(k+1)} a_{i-k-q(k+1)}}^{\iota,\, u_{i,v,q}:\, w_1}.\end{aligned}$$

Without loss of generality, it is assumed that the subwords $a_1 \cdots a_n$ at the beginning of the word of type ι and at the beginning of the word of type ζ coincide. We have

$$\begin{aligned} w_1 &= a_1^2(a_2 \cdots a_n)^2 v' b_{i+q'(k+1)+1} \cdots b_{i+q'(k+1)+1-q'(k+1)} b_{i+1-(k+1)} \cdots b_{i+1-q(k+1)}, \\ w_2 &= a_1 \cdots a_n v' a_{i+q'(k+1)+1+k} c_{i+q'(k+1)+1+k} \cdots c_{i+q'(k+1)+1+k-q'(k+1)} \cdot \\ &\qquad \cdot c_i \cdots c_{i-q(k+1)} a_{i-k-q(k+1)}. \end{aligned}$$

We have

$$\begin{aligned} w_1 &= a_1^2(a_2 \cdots a_n)^2 v' b_{i+q'(k+1)+1} \cdots b_{i+q'(k+1)+1-q'(k+1)} b_{i+1-(k+1)} \cdots b_{i+1-q(k+1)} \\ &= a_1^2(a_2 \cdots a_n)^2 v' b_{i+(q'+1)(k+1)+1-(k+1)} \cdots b_{i+(q'+1)(k+1)+1-(q'+1)(k+1)} \cdots \\ &\qquad \cdots b_{i+(q'+1)(k+1)+1-(q'+1+q)(k+1)}. \end{aligned}$$

Applying $u_{i+(q'+1)(k+1),v',q'+1+q}$, we get

$$\begin{aligned} w_2 &= a_1 \cdots a_n v' a_{i+q'(k+1)+1+k} c_{i+q'(k+1)+1+k} \cdots c_{i+q'(k+1)+1+k-q'(k+1)} c_i \cdot \\ &\qquad \cdot c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \\ &= a_1 \cdots a_n v' a_{i+(q'+1)(k+1)} c_{i+(q'+1)(k+1)} \cdots c_{i+(q'+1)(k+1)-q'(k+1)} \cdots \\ &\qquad \cdots c_{i+(q'+1)(k+1)-(q'+1+q)(k+1)} a_{i-k-q(k+1)} \\ &\succ a_1^2(a_2 \cdots a_n)^2 v' b_{i+(q'+1)(k+1)+1-(k+1)} \cdots b_{i+(q'+1)(k+1)+1-(q'+1)(k+1)} \cdots \\ &\qquad \cdots b_{i+(q'+1)(k+1)+1-(q'+1+q)(k+1)} \\ &= w_1. \end{aligned}$$

Case 22: Overlaps of the form (θ, ζ), $[(\iota, \eta)]$. We apply $d_{i,v,q}$ to the subword of type ζ to get w_1 and we apply $s_{i',v',q'}$ to the subword of type θ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. Let $w_0 = a_1 \cdots a_n$. We have

$$\begin{aligned} w &= a_1 \cdots a_n v b_i \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \\ &= \overbrace{\underbrace{w_0 v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_{i+1} \cdots a_{i'-1-q'(k+1)}}_{\theta,\ s_{i',v',q'}:\ w_2} \cdots a_{i-2} b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}}^{\zeta,\ d_{i,v,q}:\ w_1}. \end{aligned}$$

Without loss of generality, it is assumed that the subwords $w_0 = a_1 \cdots a_n$ at the beginning of the word of type ζ and at the beginning of the word of type θ coincide. We have

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_i a_{i+k} c_{i+k} \cdots c_{i+k-q(k+1)}, \\ w_2 &= a_1^2(a_2 \cdots a_n)^2 v' c_{i'+k-(k+1)} \cdots c_{i'+k-q'(k+1)} a_{i'-q'(k+1)} \cdots a_{i-2} \cdot \\ &\qquad \cdot b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)}. \end{aligned}$$

Applying h_i and and a sequence of transformations r, we get

$$\begin{aligned} w_1 &= a_1 \cdots a_n v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_i a_{i+k} c_{i+k} \cdots c_{i+k-q(k+1)} \\ &\succ a_1 \cdots a_n v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_{i-1} a_1 \cdots a_n \cdot \\ &\qquad \cdot c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} \\ &\succ \ldots \succ a_1^2(a_1 \cdots a_n)^2 v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_{i-1} \cdot \\ &\qquad \cdot c_{i+k-(k+1)} \cdots c_{i+k-q(k+1)} \\ &= a_1^2(a_1 \cdots a_n)^2 v' a_{i'} b_{i'} \cdots b_{i'-(q'-1)(k+1)} a_{i'+1-q'(k+1)} \cdots a_{i-1} c_{i-1} \cdots c_{i-1-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

We have

$$\begin{aligned} w_2 &= a_1^2(a_2\cdots a_n)^2v'c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)}a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\ &\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2v'c_{i'-1}\cdots c_{i'-1-(q'-1)(k+1)}a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\ &\qquad \cdot b_{i-k-1}\cdots b_{i-k-1-(q-1)(k+1)}a_{i-1-q(k+1)}. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2 \geq w_3$.

Case 23: Overlaps of the form (θ, η), $[(\iota, \zeta)]$. We apply $e_{i,v,q}$ to the subword of type η to get w_1 and we apply $s_{i+(q'+1)(k+1),v',q'}$ to the subword of type θ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. Let $w_0 = a_1\cdots a_n$. We have

$$\begin{aligned} w &= w_0vc_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \\ &= \overbrace{\underbrace{w_0v'a_{i+k+1+q'(k+1)}b_{i+k+1+q'(k+1)}\cdots b_{i+k+1+q'(k+1)-q'(k+1)}a_{i+k}}_{\theta,\ s_{i+(q'+1)(k+1),v',q'}:\ w_2}\cdots a_{i-2k}c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}\cdot}^{\eta,\ e_{i,v,q}:\ w_1} \end{aligned}$$

Without loss of generality, it is assumed that the subwords $w_0 = a_1\cdots a_n$ at the beginning of the word of type η and at the beginning of the word of type θ coincide. We have

$$\begin{aligned} w_1 &= a_1\cdots a_nv'a_{i+k+1+q'(k+1)}b_{i+k+1+q'(k+1)}\cdots b_{i+k+1+q'(k+1)-q'(k+1)}\cdot \\ &\qquad \cdot a_{i+1}b_{i+1}\cdots b_{i+1-q(k+1)}, \\ w_2 &= a_1^2(a_2\cdots a_n)^2v'c_{i+k+1+q'(k+1)+k-(k+1)}\cdots c_{i+k+1+q'(k+1)+k-q'(k+1)}a_{i+2k}\cdots a_{i-2k}\cdot \\ &\qquad \cdot c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}. \end{aligned}$$

Applying $s_{(q'+1)(k+1),v',q'+1}$, we get

$$\begin{aligned} w_1 &= a_1\cdots a_nv'a_{i+k+1+q'(k+1)}b_{i+k+1+q'(k+1)}\cdots b_{i+k+1+q'(k+1)-q'(k+1)}\cdot \\ &\qquad \cdot a_{i+1}b_{i+1}\cdots b_{i+1-q(k+1)} \\ &= a_1\cdots a_nv'a_{i+(q'+1)(k+1)}b_{i+(q'+1)(k+1)}\cdots b_{i+(q'+1)(k+1)-q'(k+1)}\cdot \\ &\qquad \cdot b_{i+(q'+1)(k+1)-(q'+1)(k+1)}a_{i-1}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\ &\succ a_1^2(a_2\cdots a_n)^2v'c_{i+(q'+1)(k+1)+k-(k+1)}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}\cdot \\ &\qquad \cdot c_{i+(q'+1)(k+1)+k-(q'+1)(k+1)}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2v'c_{i+(q'+1)(k+1)+k-(k+1)}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}a_{i+2k}\cdots a_{i-2k}\cdot \\ &\qquad \cdot a_{i-k}b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2v'c_{i+(q'+1)(k+1)+k-(k+1)}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}a_{i+2k}\cdots a_{i-2k}\cdot \\ &\qquad \cdot a_{i-k}b_{i-k}\cdots b_{i-k-(q-1)(k+1)} \\ &= w_1'. \end{aligned}$$

We have

$$\begin{aligned} w_2 &= a_1^2(a_2\cdots a_n)^2v'c_{i+(q'+1)(k+1)+k-(k+1)}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}a_{i+2k}\cdots a_{i-2k}\cdot \\ &\qquad \cdot c_{i-(k+1)}\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2v'c_{i+(q'+1)(k+1)+k-(k+1)}\cdots c_{i+(q'+1)(k+1)+k-q'(k+1)}a_{i+2k}\cdots a_{i-2k}\cdot \\ &\qquad \cdot c_{i-k-1}\cdots c_{i-k-1-(q-1)(k+1)}a_{i-k-1-k-(q-1)(k+1)}. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1', w_2 \geq w_3$.

Case 24 : Overlaps of the form (θ,θ), $[(\iota,\iota)]$. We apply $s_{i,v,q}$ to a subword of type θ to get w_1, and we apply $s_{i',v',q'}$ to the other subword of type θ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. Let $w_0 = a_1\cdots a_n$. We have

$$\begin{aligned} w &= a_1\cdots a_n v a_i b_i\cdots b_{i-q(k+1)}a_{i-1-q(k+1)} \\ &= \overbrace{\underbrace{w_0 v' a_{i'}b_{i'}\cdots b_{i'-(q'-1)(k+1)}a_{i'+1-q'(k+1)}\cdots a_i\cdots a_{i'-1-q'(k+1)}}_{\theta,\ s_{i',v',q'}:\ w_2}\cdots a_{i-2}b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}}^{\theta,\ s_{i,v,q}:\ w_1}. \end{aligned}$$

Without loss of generality, it is assumed that the subwords $w_0 = a_1\cdots a_n$ at the beginning of the two words of type θ coincide. We have

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}b_{i'}\cdots b_{i'-(q'-1)(k+1)}a_{i'+1-q'(k+1)}\cdots a_{i-1}\cdot \\ &\qquad \cdot c_{i+k-(k+1)}c_{i+k-2(k+1)}\cdots c_{i+k-q(k+1)}, \\ w_2 &= a_1^2(a_2\cdots a_n)^2 v' c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)}a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\ &\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)}. \end{aligned}$$

We have

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}b_{i'}\cdots b_{i'-(q'-1)(k+1)}a_{i'+1-q'(k+1)}\cdots a_{i-1}\cdot \\ &\qquad \cdot c_{i+k-(k+1)}\cdots c_{i+k-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' a_{i'}b_{i'}\cdots b_{i'-(q'-1)(k+1)}a_{i'+1-q'(k+1)}\cdots a_{i-1}\cdot \\ &\qquad \cdot c_{i-1}\cdots c_{i-1-(q-1)(k+1)}, \\ w_2 &= a_1^2(a_2\cdots a_n)^2 v' c_{i'+k-(k+1)}\cdots c_{i'+k-q'(k+1)}a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\ &\qquad \cdot b_{i-(k+1)}\cdots b_{i-q(k+1)}a_{i-1-q(k+1)} \\ &= a_1^2(a_2\cdots a_n)^2 v' c_{i'-1}\cdots c_{i'-1-(q'-1)(k+1)}a_{i'-q'(k+1)}\cdots a_{i-2}\cdot \\ &\qquad \cdot b_{i-k-1}\cdots b_{i-k-1-(q-1)(k+1)}a_{i-k-1-1-(q-1)(k+1)}. \end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2 \geq w_3$.

Case 25: Overlaps of the form (θ,ι), $[(\iota,\theta)]$. We apply $u_{i,v,q}$ to the subword of type ι to get w_1, and we apply $s_{i+1+q'(k+1),v',q'}$ to the subword of type θ to get w_2. There are two kinds of overlap. We consider the first kind of overlap, the second kind, as in *case 10*, can be solved similarly. We have

$$\begin{aligned} w &= a_1\cdots a_n v a_i c_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \\ &= \overbrace{\underbrace{a_1\cdots a_n v' a_{i+1+q'(k+1)}b_{i+1+q'(k+1)}\cdots b_{i+1+q'(k+1)-q'(k+1)}a_i}_{\theta,\ s_{i+1+q'(k+1),v',q'}:\ w_2}c_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}}^{\iota,\ u_{i,v,q}:\ w_1}. \end{aligned}$$

Without loss of generality, it is assumed that the subwords $a_1\cdots a_n$ at the beginning of the word of type ι and at the beginning of the word of type θ coincide. We have

$$\begin{aligned} w_1 &= a_1^2(a_2\cdots a_n)^2 v' a_{i+1+q'(k+1)}b_{i+1+q'(k+1)}\cdots b_{i+1+q'(k+1)-q'(k+1)}\cdot \\ &\qquad \cdot b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)}, \\ w_2 &= a_1^2(a_2\cdots a_n)^2 v' c_{i+1+q'(k+1)+k-(k+1)}\cdots c_{i+1+q'(k+1)+k-q'(k+1)}\cdot \\ &\qquad \cdot c_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)}. \end{aligned}$$

We have

$$\begin{aligned}
w_1 &= a_1^2(a_2\cdots a_n)^2v'a_{i+1+q'(k+1)}b_{i+1+q'(k+1)}\cdots b_{i+1+q'(k+1)-q'(k+1)}\cdot \\
&\qquad\qquad \cdot b_{i+1-(k+1)}\cdots b_{i+1-q(k+1)} \\
&= a_1^2(a_2\cdots a_n)^2v'a_{i+1+q'(k+1)}b_{i+1+q'(k+1)}\cdots b_{i+1+q'(k+1)-q'(k+1)}\cdot \\
&\qquad\qquad \cdot b_{i+1+q'(k+1)-(q'+1)(k+1)}\cdots b_{i+1+q'(k+1)-(q'+q)(k+1)}, \\
w_2 &= a_1^2(a_2\cdots a_n)^2v'c_{i+1+q'(k+1)+k-(k+1)}\cdots c_{i+1+q'(k+1)+k-q'(k+1)}\cdot \\
&\qquad\qquad \cdot c_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \\
&= a_1^2(a_2\cdots a_n)^2v'c_{i+q'(k+1)}\cdots c_{i+q'(k+1)-(q'-1)(k+1)}c_i\cdots c_{i-q(k+1)}a_{i-k-q(k+1)} \\
&= a_1^2(a_2\cdots a_n)^2v'c_{i+q'(k+1)}\cdots c_{i+q'(k+1)-(q'-1)(k+1)}\cdot \\
&\qquad\qquad \cdot c_{i+q'(k+1)-q'(k+1)}\cdots c_{i+q'(k+1)-(q'+q)(k+1)}a_{i+q'(k+1)-k-(q'+q)(k+1)}.
\end{aligned}$$

By Lemma 2.2, there exists $w_3 \in \mathrm{FM}_n$ such that $w_1, w_2 \geq w_3$. ∎

Note that, using the notation in the last part of the proof of the above theorem, $S_n \setminus a_1a_2\cdots a_nS_n = \{a \in S_n \mid \mathrm{card}(\pi^{-1}(a)) = 1\}$. So, elements in this set have a unique presentation in S_n. In particular, we have a natural isomorphism $S_n/(a_1\cdots a_nS_n) \cong \mathrm{FM}_n/\pi^{-1}(a_1\cdots a_nS_n)$ (again, we use the same notation for the generators of FM_n and those of S_n).

3 The algebra $K[S_n(H)]$ defined by dihedral type relations

Recall that if R is a finitely generated algebra over a field K, with a set of generators $a_1, \ldots, a_n$, and $\pi\colon K\langle x_1, \ldots, x_n\rangle \longrightarrow R$ is the unique homomorphism from the free algebra $K\langle x_1, \ldots, x_n\rangle$ to R such that $\pi(x_i) = a_i$, for all i, and we have a fixed order on the free monoid $\mathrm{FM}_n = \langle x_1, \ldots, x_n\rangle$ compatible with the multiplication in FM_n, then R is an automaton algebra with respect to the presentation π and the fixed order on FM_n if and only if the ideal I of FM_n consisting of all leading monomials of elements of $\ker(\pi)$ is a regular language. A set of elements of FM_n is a regular language if it is obtained from a finite subset of FM_n by applying a finite number of operations of union, multiplication and operation $*$, defined by $T^* = \bigcup_{i\geq 1} T^i$, for $T \subseteq \mathrm{FM}_n$. Multiplication means multiplication of non-empty subsets of FM_n: if A and B are non-empty subsets of FM_n, then $AB = \{ab \mid a \in A \text{ and } b \in B\}$. For example, $\{x_1, x_2\}x_3^* = \{x_1x_3^i \mid i \geq 1\} \cup \{x_2x_3^i \mid i \geq 1\}$ and this is a regular language. Note that FM_n is a regular language because $\mathrm{FM}_n = \{1, x_1, x_2, \ldots, x_n\}^*$. The set $N(R) = \mathrm{FM}_n \setminus I$ is called the set of normal words of R, with respect to the given presentation of R and the fixed order on FM_n. By [16, Lemma of page 96], I is a regular language if and only if $N(R)$ is a regular language.

Let H be a subgroup of Sym_n that contains $\langle(1, 2, \ldots, n)\rangle$ as a normal subgroup of index 2, where $n > 3$. We denote $S_n(H)$ by S_n. An easy consequence of Theorem 2.3 is the following result.

Theorem 3.1 *Let K be a field. Then $K[S_n]$ is an automaton algebra with respect to the given presentation of S_n and the length-lexicographical order $\ll$ on FM_n generated by $a_1 \ll a_2 \ll \cdots \ll a_n$. (Here we use the same notation for the generators of FM_n and those of S_n).*

Proof. For $w \in \mathrm{FM}_n$, we define $w^* = \{w^i \mid i \geq 0\}$. By Theorem 2.3, the set of normal words of the

algebra $K[S_n]$ is

$$\begin{aligned} N(K[S_n]) = a_1^*(\mathrm{FM}_n \setminus (I \cup a_2 \cdots a_n \,\mathrm{FM}_n)) \cup (a_2 \cdots a_n)^*(\mathrm{FM}_n \setminus (I \cup a_1 \,\mathrm{FM}_n)) \cup \\ \cup\ a_1 a_1^*(a_2 \cdots a_n)(a_2 \cdots a_n)^* \cdot \\ \cdot (\mathrm{FM}_n \setminus (I \cup a_1 \,\mathrm{FM}_n \cup a_2 \cdots a_n \,\mathrm{FM}_n \cup I_\zeta \cup I_\eta \cup I_\theta \cup I_\iota)), \end{aligned}$$

where $I = \bigcup_{\sigma \in H} \mathrm{FM}_n \, a_{\sigma(1)} \cdots a_{\sigma(n)} \, \mathrm{FM}_n$,

$$\begin{aligned} I_\zeta &= \textstyle\bigcup_{q=0}^{n-1} \bigcup_{i=n-k+1}^{n-1} \mathrm{FM}_n (b_i b_{i-(k+1)} \cdots b_{i-(n-1)(k+1)})^* \cdot \\ &\qquad \cdot b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \, \mathrm{FM}_n, \\ I_\eta &= \textstyle\bigcup_{q=0}^{n-1} \bigcup_{i=0}^{n-k} \mathrm{FM}_n (c_i c_{i-(k+1)} \cdots c_{i-(n-1)(k+1)})^* \cdot \\ &\qquad \cdot c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \, \mathrm{FM}_n, \\ I_\theta &= \textstyle\bigcup_{q=0}^{n-1} \bigcup_{i=1}^{n} \mathrm{FM}_n \, a_i (b_i b_{i-(k+1)} \cdots b_{i-(n-1)(k+1)})^* \cdot \\ &\qquad \cdot b_i b_{i-(k+1)} \cdots b_{i-q(k+1)} a_{i-1-q(k+1)} \, \mathrm{FM}_n, \\ I_\iota &= \textstyle\bigcup_{q=0}^{n-1} \bigcup_{i=1}^{n} \mathrm{FM}_n \, a_i (c_i c_{i-(k+1)} \cdots c_{i-(n-1)(k+1)})^* \cdot \\ &\qquad \cdot c_i c_{i-(k+1)} \cdots c_{i-q(k+1)} a_{i-k-q(k+1)} \, \mathrm{FM}_n \, . \end{aligned}$$

By [16, Lemma of page 96], $N(K[S_n])$ is a regular language. Therefore $K[S_n]$ is an automaton algebra. ∎

It follows from the beginning of Section 2, that the universal group of S_n, which means the group with the same presentation as S_n, is

$$\begin{aligned} G_n = \mathrm{gr}(a_1, a_2, \ldots, a_n \mid a_1 a_2 \cdots a_{n-1} a_n &= a_i a_{i+1} a_{i+2} \cdots a_{i-2} a_{i-1} \\ &= a_i a_{i+k} a_{i+2k} \cdots a_{i-2k} a_{i-k},\ 1 \le i \le n), \end{aligned}$$

where $k^2 \equiv 1 \mod n$.

Theorem 3.2 *The universal group G_n of $S_n(H)$ is a unique product group.*

Proof. Consider the groups

$$H_n = \mathrm{gr}(a_2, \ldots, a_n \mid a_2 a_3 \cdots a_n = a_{1+k} a_{1+2k} \cdots a_{1+(n-1)k}),$$

and

$$T_n = \mathrm{gr}(z) \times H_n.$$

Note that $(z, 1)$ is a central element of T_n and

$$\begin{aligned} (z, (a_2 a_3 \cdots a_n)^{-1})(1, a_2)(1, a_3) \cdots (1, a_n) &= (z, 1) \\ \text{and} \quad (z, (a_2 a_3 \cdots a_n)^{-1})(1, a_{1+k})(1, a_{1+2k}) \cdots (1, a_{1+(n-1)k}) &= (z, 1) \end{aligned}$$

Hence there exists a unique homomorphism $f : G_n \to T_n$ such that

$$\begin{aligned} f(a_i) &= (1, a_i) \quad \text{for } i = 2, \ldots, n, \text{ and} \\ f(a_1) &= (z, (a_2 \cdots a_n)^{-1}). \end{aligned}$$

Note that $a_1a_2\cdots a_n$ is a central element in G_n (see page 2) and that $a_2a_3\cdots a_n = a_{1+k}a_{1+2k}\cdots a_{1+(n-1)k}$ in G_n. Furthermore, because the defining relations of G_n are homogeneous with respect to the degree of each a_i, the subgroup of G_n generated by $a_2,\ldots,a_n$ intersects trivially with the subgroup generated by $a_1a_2\cdots a_n$. Hence there exists a unique homomorphism $g : T_n \to G_n$ such that

$$\begin{aligned} g(1,a_i) &= a_i \quad \text{for } i = 2,\ldots,n \text{ and} \\ g(z,1) &= a_1a_2\cdots a_n \end{aligned}$$

It is easy to check that $g = f^{-1}$. Hence

$$G_n \cong T_n.$$

Recall that a group G is said to be locally indicable if every non-trivial finitely generated subgroup of G can be mapped homomorphically onto a finite cyclic group. Such groups are right ordered [13, Exercise 9, page 638] and thus t.u.p. groups. Indicable groups were first introduced by G. Higman in [9]. It is well known that the group algebra of a u.p. group only has trivial units (see [13, Lemma 13.1.9]) and thus no zero divisors.

By a result of Brodskiĭ [2], every torsion free one-relator group is locally indicable (see [10, Theorem 3.1] for a short proof). Since the word $a_2a_3\cdots a_na_{1+(n-1)k}^{-1}a_{1+(n-2)k}^{-1}\cdots a_{1+k}^{-1}$ is not a proper power, by [14, Theorem 4.12], H_n is torsion free. Hence H_n is locally indicable, thus it is a unique product group. A direct product of u.p. groups is u.p, thus G_n is a u.p. group. ∎

Note that the monoid $S_n = S_n(H)$ is not cancellative. Thus S_n is not embedded in G_n. Let $z = a_1a_2\ldots a_n \in S_n$, a central element in S_n.

Lemma 3.3 *The subsemigroup zS_n of S_n is cancellative and $G_n = zS_n\langle z\rangle^{-1}$.*

Proof. Since z is central, it follows from the defining relations of S_n that for every $w \in S_n$ there exists $u \in S_n$ such that $uw = z^l$ for some positive integer l. Let $w_k = a_1^{i_k}(a_1a_2\cdots a_n)(a_2\cdots a_n)^{j_k}b_k, k = 1,2$, be two elements of zS_n written in the canonical form established in Theorem 2.3. Suppose that $ww_1 = ww_2$, for some $w \in zS_n$. Then $z^lw_1 = z^lw_2$, for some positive integer l. But the canonical form of z^lw_k is

$$z^lw_k = a_1^{i_k+l}(a_1a_2\cdots a_n)(a_2\cdots a_n)^{j_k+l}b_k,$$

for $k = 1,2$. Hence $w_1 = w_2$ and therefore, zS_n is left cancellative. Similarly one can prove that zS_n is right cancellative. Therefore, it follows that zS_n is cancellative and that the central localization $zS_n\langle z\rangle^{-1}$ is the group of fractions of zS_n. Now it is easy to see that $G_n = zS_n\langle z\rangle^{-1}$. ∎

In [5, Section 2] some general results on the structure of the algebra $K[S_n(H)]$ are proved (for arbitrary subgroups H of the symmetric group of degree n). In particular, if $z = a_1a_2\cdots a_n \in S_n(H)$ is central, then $\mathcal{J}(K[S_n(H)] \subseteq K[zS_n(H)]$ [5, Proposition 2.6]. Using this in the present case, together with the fact that zS_n is cancellative, we obtain the following result.

Theorem 3.4 *Let K be a field. Then $\mathcal{J}(K[S_n]) = \{0\}$.*

Proof. We know that G_n is a u.p. group. Hence $K[G_n]$ has only trivial units ([15] and [13, Lemma 13.1.9]). Let $\alpha \in \mathcal{J}(K[S_n])$. By [5, Proposition 2.6], $\alpha \in K[zS_n]$. There exists $\beta \in \mathcal{J}(K[S_n])$ such that

$$\alpha + \beta + \alpha\beta = 0.$$

We also have that $\beta \in K[zS_n]$. Since

$$(1+\alpha)(1+\beta) = 1,$$

and $K[G_n]$ has only trivial units, there exist $\lambda_1, \lambda_2 \in K$ and $w_1, w_2 \in zS_n \cup \{1\}$, such that

$$1 + \alpha = \lambda_1 w_1 \quad \text{and} \quad 1 + \beta = \lambda_2 w_2$$

and $\lambda_1\lambda_2 w_1 w_2 = 1$. Hence $w_1 w_2 = 1$. Because S_n only has trivial units, we obtain that $w_1 = w_2 = 1$ and

$$\alpha = \lambda_1 w_1 - 1 = (\lambda_1 - 1) \cdot 1.$$

Hence $\lambda_1 = 1$, and $\alpha = 0$. ∎

The proof of the following two results is as the proof of Lemma 2.5 and Proposition 2.6 of [4].

Lemma 3.5 *$P = a_1 a_2 \cdots a_n S_n$ is a prime ideal of S_n. Moreover, every prime ideal Q of $K[S_n]$ such that $Q \cap S_n \neq \emptyset$ contains P. Furthermore, $K[P]$ is a height one prime ideal of $K[S_n]$.*

Note that if Q is a prime ideal of S_n, then $K[Q]$ is a prime ideal of $K[S_n]$. Indeed, by the proof of Lemma 3.5, $P \subseteq Q$ and, hence, by [11, Proposition 24.2], $K[S_n]/K[Q]$ is a prime monomial algebra. In particular, since $n > 2$, the submonoid $\langle a_1, a_2 \rangle$ of S_n/P is a free monoid of rank two. Therefore $K[S_n]/K[P]$ is not a PI-algebra. Hence, by a recent result of Okniński [12, Theorem 0.1], and by a result of Bell and Colak [1, Theorem 1.2], we obtain the following result.

Proposition 3.6 *Assume that Q is a prime ideal of S_n. Then $K[S_n]/K[Q]$ is either a prime PI-algebra or a (left and right) primitive ring or it has a nonzero locally nilpotent Jacobson radical.*

Assume, furthermore, that Q is finitely generated. Then $K[S_n]/K[Q]$ is either a prime PI-algebra or a (left and right) primitive ring. In particular, $K[S_n]/K[P]$ is a (left and right) primitive ring.

References

[1] J.P. Bell and P. Colak, Primitivity of finitely presented monomial algebras, J. Pure Appl. Algebra 213 (2009), 1299–1305.

[2] S. D. Brodskiĭ Equations over groups and groups with one defining relation, Sibirsk. Mat. Zh. 25(2) (1984), 84–103; Siberian Math. J. 25(2) (1984), 235–251.

[3] F. Cedó, E. Jespers and J. Okniński, The radical of the four generated algebra of the alternating type, Contemporary Math 499 (2009), 1–26.

[4] F. Cedó, E. Jespers and J. Okniński, Finitely presented algebras and groups defined by permutation relations, J. Pure App. Algebra 214 (2010), 1095–1102.

[5] F. Cedó, E. Jespers and J. Okniński, Algebras and groups defined by permutation relations of alternating type, J. Algebra 324 (2010) 1290–1313.

[6] F. Cedó, E. Jespers and G. Klein, Finitely presented monoids and algebras defined by permutation relations of abelian type, J. Pure App. Algebra 216 (2012) 1033–1039.

[7] F. Cedó, E. Jespers and G. Klein, Construction of a two unique product semigroup defined by permutation relations of quaternion type, in preparation.

[8] P.M. Cohn, Universal Algebra, revised edition, D. Reidel Publishing Company, Dordrecht, Holland, 1981.

[9] G. Higman, The units of group-rings, Proceedings of the London Mathematical Society, Second Series, 46 (1940), 231–248.

[10] J. Howie, A Short proof of a theorem of Brodskii, Publicacions matematiques. Bellaterra : Servei de Publicacions de la Universitat Autònoma de Barcelona, V. 44 N. 2 (2000), p. 641–647.

[11] J. Okniński, Semigroup Algebras, Marcel Dekker, New York, 1991.

[12] J. Okniński, Trichotomy for finitely generated monomial algebras, J. Algebra 417 (2014), 145–147.

[13] D.S. Passman, The Algebraic Structure of Group Rings, Wiley, New York, 1977.

[14] W. Magnus, A. Karrass and D. Solitar, Combinatorial group theory: Presentations of groups in terms of generators and relations, Sec. Rev. Ed., Dover Publications, Inc., New York, 1976.

[15] A. Strojnowski, A note on u.p. groups, Comm. Algebra, V. 8 N. 3 (1980), 231–234.

[16] V.A. Ufnarovskiĭ, Combinatorial and Asymptotic Methods in Algebra, in: Encyclopedia of Mathematical Sciences vol. 57, pp.1–196, Springer, 1995.

F. Cedó
Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193 Bellaterra (Barcelona), Spain
cedo@mat.uab.cat

E. Jespers
Department of Mathematics
Vrije Universiteit Brussel
Pleinlaan 2, 1050 Brussel, Belgium
efjesper@vub.ac.be

G. Klein
Department of Mathematics
Vrije Universiteit Brussel
Pleinlaan 2, 1050 Brussel, Belgium
gklein@vub.ac.be